

KEK-TH-484
 KEK Preprint 97-3
 OU-HET 261
 TU-519
 April 1997

$b \rightarrow s \ell \bar{\ell}$ process and multi-Higgs doublet model

Yasuhiro Okada¹
 Yasuhiro Shimizu²,
 and Minoru Tanaka³

¹, ²*Theory Group, KEK, Tsukuba, Ibaraki, 305 Japan*

²*Department of Physics, Tohoku University
 Sendai 980-777 Japan*

³*Department of Physics, Osaka University
 Toyonaka, Osaka, 560 Japan*

Abstract

Rare b decay processes are analyzed in the multi-Higgs doublet model. Taking account of the constraint from the $b \rightarrow s \gamma$ process, the branching ratio and the forward-backward asymmetry of the final leptons for the $b \rightarrow s \ell^+ \ell^-$ process are calculated. It is shown that the branching ratio can be a few times larger than the standard model prediction and the asymmetry can be significantly different from that in the standard model. Combining these observable quantities it is possible to determine complex coupling constants associated with the charged Higgs mixing matrix. The CP violating charge asymmetry in the $b \rightarrow s \ell^+ \ell^-$ process and the branching ratio of the $b \rightarrow s \nu \bar{\nu}$ process are also calculated.

In search for new physics beyond the Standard Model (SM), flavor changing neutral current (FCNC) processes can play an important role. Since the source of the flavor changing processes only lies in the Cabbibo-Kobayashi-Maskawa (CKM) matrix in the SM, it is possible to make precise predictions for various observable quantities in FCNC processes within the SM. An important example is the branching ratio of the $b \rightarrow s \gamma$ process, which was reported to be $\text{Br}(b \rightarrow s \gamma) = (2.32 \pm 0.57 \pm 0.35) \times 10^{-4}$ from the CLEO experiment [1]. Using the observed top quark mass, the measured branching ratio is consistent with the SM prediction. Therefore this process becomes a very strong constraint on models beyond the SM such as two Higgs doublet model [2] and supersymmetric extensions of the SM [3]. In addition to the $b \rightarrow s \gamma$ process, the $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ processes can be important constraints on new physics. Current experimental bounds for these processes are only by one order of magnitudes above the SM predictions [4, 5].

In this letter we consider possible constraints on parameters in the multi-Higgs doublet model from these rare b decay processes. When the number of the Higgs doublets is more than three we can introduce complex phases in the charged Higgs mixing matrix. In this sense the multi-Higgs doublet model is a natural extension of the SM which involves a new source of the CP violation. Present phenomenological constraints on coupling constants in this model are considered, for example, in Ref.[6]. Remarkably it is pointed out in Ref.[7] that the strongest constraint on the imaginary part of coupling constant comes from the $b \rightarrow s \gamma$ process, not from the CP violating quantities such as neutron electric dipole moment. Here we study the $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ processes in the same model. We found that within the present experimental constraints including the $b \rightarrow s \gamma$ process the branching ratio can be a few times larger than the SM predictions and the forward-backward asymmetry of the final leptons for $b \rightarrow s \ell^+ \ell^-$ process can significantly differ from the SM. Combining information from the branching ratio and the forward-backward asymmetry, it is possible to determine the complex coupling constants in future experiments. We also calculate the CP violating charge asymmetry between the $b \rightarrow s \ell^+ \ell^-$ and $\bar{b} \rightarrow \bar{s} \ell^+ \ell^-$ decays, which turns out to be a few percent in the allowed region.

We consider the multi-Higgs doublet model with the following Yukawa couplings,

$$\mathcal{L} = \bar{q}_L y_d d_R H_d + \bar{q}_L y_u u_R H_u + \bar{\ell}_L y_\ell e_R H_\ell + h.c., \quad (1)$$

where y_d, y_u, y_ℓ are 3×3 matrices. We have assumed that up-type quarks, down-type quarks and leptons have Yukawa couplings with only one Higgs doublet, H_u, H_d, H_ℓ respectively. In this way we can avoid FCNC effects at the tree level [8]. The couplings of physical charged Higgs bosons with fermions are then given by

$$\mathcal{L} = (2\sqrt{2}G_F)^{1/2} \sum_{i=1}^{n-1} (X_i \bar{u}_L V M_D d_R H_i^+ + Y_i \bar{u}_R M_U V d_L H_i^+ + Z_i \bar{\nu}_L M_E e_R H_i^+), \quad (2)$$

where n is the number of Higgs doublets, $H_i^+ (i = 1 \sim n - 1)$ represent mass eigenstates of charged Higgs bosons and V is the CKM matrix. New CP violating complex phases arise in the charged Higgs mixing matrix if there are three or more Higgs doublets. In such cases the coupling constants X_i, Y_i, Z_i are in general complex numbers. These are several relations among X_i, Y_i, Z_i from the requirement of unitarity of the mixing matrix [6].

The charged Higgs interactions in Eq.(2) can induce extra contribution to FCNC processes in this model. Here, we are interested in FCNC processes related to the b quark. Inclusive branching ratios of $b \rightarrow s \gamma$, $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ are calculated through the following weak effective Hamiltonian at the bottom scale as described in [9, 10];

$$\mathcal{H} = -\frac{4G_F}{\sqrt{2}} V_{tb} V_{ts}^* \sum_{i=1}^{10} C_i O_i. \quad (3)$$

The relevant operators at the bottom scale are

$$O_7 = \frac{e}{16\pi^2} m_b \bar{s}_L \sigma_{\mu\nu} b_R F^{\mu\nu}, \quad (4)$$

$$O_8 = \frac{g_s}{16\pi^2} m_b \bar{s}_L T^a \sigma_{\mu\nu} b_R G^{a\mu\nu}, \quad (5)$$

$$O_9 = \frac{e^2}{16\pi^2} \bar{s}_L \gamma^\mu b_L \bar{\ell} \gamma_\mu \ell, \quad (6)$$

$$O_{10} = \frac{e^2}{16\pi^2} \bar{s}_L \gamma^\mu b_L \bar{\ell} \gamma_\mu \gamma^5 \ell, \quad (7)$$

$$O_{11} = \frac{e^2}{16\pi^2 \sin^2 \theta_W} \bar{s}_L \gamma^\mu b_L \sum_{i=e,\mu,\tau} \bar{\nu}_i \gamma_\mu (1 - \gamma^5) \nu_i, \quad (8)$$

where e , g_s are QED and strong coupling constants respectively, and θ_W is the weak mixing angle. Using renormalization group equations of QCD the Wilson coefficients C_i 's at the bottom scale are related to those at the weak scale. New physics effects enter through the Wilson coefficients at the weak scale. From the charged Higgs interactions these coefficients receive the following new contributions,

$$C_7^H = -\frac{1}{2} \sum_{i=1}^{n-1} x_{th_i} \left[|Y_i|^2 \left(\frac{2}{3} F_1(x_{th_i}) + F_2(x_{th_i}) \right) + X_i Y_i^* \left(\frac{2}{3} F_3(x_{th_i}) + F_4(x_{th_i}) \right) \right], \quad (9)$$

$$C_8^H = -\frac{1}{2} \sum_{i=1}^{n-1} x_{th_i} \left[|Y_i|^2 F_1(x_{th_i}) + X_i Y_i^* F_3(x_{th_i}) \right], \quad (10)$$

$$C_9^H = -D^H + \frac{1 - 4 \sin^2 \theta_W}{\sin^2 \theta_W} C^H, \quad (11)$$

$$C_{10}^H = -\frac{C^H}{\sin^2 \theta_W}, \quad (12)$$

$$C_{11}^H = -C^H. \quad (13)$$

Here C^H and D^H are given by

$$C^H = \sum_{i=1}^{n-1} \frac{1}{8} |Y_i|^2 x_{th_i} x_{tW} (F_3(x_{th_i}) + F_4(x_{th_i})), \quad (14)$$

$$D^H = \sum_{i=1}^{n-1} x_{th_i} |Y_i|^2 \left(\frac{2}{3} F_5(x_{th_i}) - F_6(x_{th_i}) \right), \quad (15)$$

where $x_{tW} = m_t^2/m_W^2$, $x_{th_i} = m_t^2/m_{H_i}^2$. The functions F_1 - F_6 are defined as follows: $F_1(x) = \frac{1}{12(x-1)^4}(x^3 - 6x^2 + 3x + 2 + 6x \ln x)$, $F_2(x) = \frac{1}{12(x-1)^4}(2x^3 + 3x^2 - 6x + 1 - 6x^2 \ln x)$, $F_3(x) = \frac{1}{2(x-1)^3}(x^2 - 4x + 3 + 2 \ln x)$, $F_4(x) = \frac{1}{2(x-1)^3}(x^2 - 1 - 2x \ln x)$, $F_5(x) = \frac{1}{36(x-1)^4}(7x^3 - 36x^2 + 45x - 16 + (18x - 12) \ln x)$, $F_6(x) = \frac{1}{36(x-1)^4}(-11x^3 + 18x^2 - 9x + 2 + 6x^3 \ln x)$. The Wilson coefficients at the weak scale are then given by

$$C_i(m_W) = C_i^{SM}(m_W) + C_i^H(m_W), \quad (16)$$

where C_i^{SM} is contribution from the SM [9, 10]. After taking account of the QCD corrections at the leading logarithmic order, $C_i(m_b)$ can be expressed as,

$$C_7(m_b) = C_7(m_W)\eta^{\frac{16}{23}} + C_8(m_W)\frac{8}{3}(\eta^{\frac{14}{23}} - \eta^{\frac{16}{23}}) + \tilde{C}_7, \quad (17)$$

$$C_8(m_b) = C_8(m_W)\eta^{\frac{14}{23}} + \tilde{C}_8, \quad (18)$$

$$C_9(m_b) = C_9(m_W) + \tilde{C}_9, \quad (19)$$

$$C_{10}(m_b) = C_{10}(m_W), \quad (20)$$

$$C_{11}(m_b) = C_{11}(m_W), \quad (21)$$

where $\eta = \alpha_s(m_W)/\alpha_s(m_b)$. $\tilde{C}_7 \sim \tilde{C}_9$ are constants which depend on the QCD coupling constant. Detailed formulas are found in [10, 11]. Numerically these are given by $\tilde{C}_7 = -0.17$, $\tilde{C}_8 = -0.077$, $\tilde{C}_9 = 1.9$ for $\alpha_s(m_Z) = 0.12$.

The $b \rightarrow s \gamma$ branching ratio is then given by

$$\text{Br}(b \rightarrow s \gamma) = \text{Br}(b \rightarrow c e \bar{\nu}) \frac{6\alpha_{em}}{\pi g(m_c/m_b)} \left| \frac{V_{ts} V_{tb}^*}{V_{cb}} \right|^2 |C_7(m_b)|^2, \quad (22)$$

where the phase space factor $g(z)$ is given by $g(z) = 1 - 8z^2 + 8z^6 - z^8 - 24z^4 \ln z$. For the $b \rightarrow s \ell^+ \ell^-$ process, the differential branching ratio and the forward-backward asymmetry of leptons in the lepton-center-of-mass frame are given by

$$\frac{d\text{Br}(b \rightarrow s \ell^+ \ell^-)}{d\hat{s}} = \text{Br}(b \rightarrow c e \bar{\nu}) \frac{\alpha_{em}^2}{4\pi} \left| \frac{V_{ts} V_{tb}^*}{V_{cb}} \right|^2 \frac{1}{g(m_c/m_b)} (1 - \hat{s})^2 \quad (23)$$

$$\times \left[(|C_9 + Y(\hat{s})|^2 + |C_{10}|^2)(1 + 2\hat{s}) + \frac{4}{\hat{s}} |C_7|^2 (2 + \hat{s})^2 + 12\text{Re}C_7^*(C_9 + Y(\hat{s})) \right], \quad (24)$$

$$A_{FB}(\hat{s}) = \frac{-3\text{Re}C_{10}^*[(C_9 + Y(\hat{s}))\hat{s} + 2C_7]}{(|C_9 + Y(\hat{s})|^2 + |C_{10}|^2)(1 + 2\hat{s}) + \frac{4}{\hat{s}} |C_7|^2 (2 + \hat{s})^2 + 12\text{Re}C_7^*(C_9 + Y(\hat{s}))},$$

where $\hat{s} = s/m_b^2 = (p_+ + p_-)^2/m_b^2$ and $p_+(p_-)$ is the four momentum of $\ell^+(\ell^-)$. $Y(\hat{s})$ represents the contribution from the charm quark loop at the m_b scale. See Ref.[11] for details. Here we neglect the J/ψ and ψ' resonance contributions. The branching ratio for the $b \rightarrow s \nu \bar{\nu}$ is given by

$$\sum_{i=e,\mu,\tau} \text{Br}(b \rightarrow s \nu_i \bar{\nu}_i) = 3 \text{Br}(b \rightarrow c e \bar{\nu}) \frac{\alpha^2}{4\pi^2 \sin^4 \theta_W} \left| \frac{V_{tb} V_{ts}^*}{V_{cb}} \right|^2 \frac{1}{g(m_c/m_b)} |C_{11}|^2. \quad (25)$$

Using the above formulas, it is now straight-forward to evaluate the branching ratios and the asymmetry numerically in the multi-Higgs doublet model. In the followings, we assume that only one of the physical charged Higgs boson is light and neglect the effects of other physical charged Higgs bosons. Then we keep only one term in the summation in the expressions for C_i^H 's. Dropping the index i for the lightest charged Higgs bosons, the relevant parameters are $|Y|^2$, XY^* and the mass of the charged Higgs boson. It should be noted that the coefficients C_7 and C_8 depend on both $|Y|^2$ and XY^* whereas C_9 and C_{10} only contain $|Y|^2$.

Since the $b \rightarrow s \gamma$ branching ratio is already observed and is consistent with the SM prediction, we use $\text{Br}(b \rightarrow s \gamma)$ to solve $\text{Im}XY^*$ in terms of $|Y|^2$ and $\text{Re}XY^*$ for each value of the charged Higgs mass. Then other observable quantities for $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ processes can be calculated as functions of two parameters. We should also take into account constraints from other processes. Besides the $b \rightarrow s \gamma$ process, the B^0 - $\bar{B}^0$ mixing and the $Z \rightarrow b \bar{b}$ process give the most strong constraints on the possible value of $|Y|^2$ [6]. The contribution to the B^0 - $\bar{B}^0$ mixing from the charged Higgs boson is expressed as

$$M_{12}^H = \frac{G_F^4}{64\pi^2} m_W^2 \eta_B (V_{td} V_{tb}^*)^2 \frac{4}{3} B_B f_B^2 m_B x_t \left[y_t I_1(y_t) |Y|^4 \right. \\ \left. + x_t (2I_2(x_t, x_H) - 8I_3(x_t, x_H) |Y|^2) \right], \quad (26)$$

where $y_t = m_t^2/m_H^2$, $x_H = m_H^2/m_W^2$, and the functions I_1 - I_3 are defined as follows.

$$I_1(x) = \frac{1+x}{(1-x)^2} + \frac{2x \ln x}{(1-x)^3}, \quad (27)$$

$$I_2(x, y) = \frac{x}{(x-y)(x-1)} + \frac{y^2 \ln(y)}{(y-1)(x-y)^2} + \frac{x(-x-xy+2y) \ln(x)}{(1-x)^2(x-y)^2}, \quad (28)$$

$$I_3(x, y) = \frac{1}{(x-y)(x-1)} + \frac{y \ln y}{(y-1)(x-y)^2} + \frac{(-x^2+y) \ln(x)}{(1-x)^2(x-y)^2}. \quad (29)$$

Here we have retained only the relevant terms which are proportional to $|Y|^4$ and $|Y|^2$. Since this quantity depends on the CKM matrix element V_{td} which has not been known well, the constraint from B^0 - $\bar{B}^0$ mixing is not very strong. Using the constraint on V_{td} from the charmless b decay ($0.005 < |V_{td}| < 0.012$) and taking account of uncertainties from f_B and B_B ($f_B \sqrt{B_B} = 200 \pm 40 \text{ MeV}$) we can deduce

$|Y| \lesssim 1.5(2.1)$ for $m_H = 100(300)$ GeV *. Here and in the followings we fix the top quark mass as 175 GeV. For the $Z \rightarrow b\bar{b}$ process we calculated the charged Higgs contribution to $R_b = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ following Ref.[12]. We define the deviation from the SM contribution ($R_b^{SM} = 0.2158$) by $\delta R_b = R_b - R_b^{SM}$. The charged Higgs contribution to δR_b is shown in Fig.1 as a function of $|Y|^2$. In this figure we also show the 3σ lower bound of δR_b from the world average value $R_b = 0.2178 \pm 0.0011$ [13]. From Fig.1 we can derive $|Y| \lesssim 0.9(1.3)$ for $m_H = 100(300)$ GeV. Note that if we use the value $R_b = 0.2159 \pm 0.0009(\text{stat}) \pm 0.0011(\text{syst})$ reported by ALEPH [14] the lower bound of δR_b shifts to -0.0041 , which corresponds to the upper bound $|Y| \lesssim 1.6(2.3)$ for $m_H = 100(300)$ GeV. Since experimental situation is not conclusive, to be conservative, we present the result in a rather wider range of $|Y|$.

Fig.2(a) shows that the ratio of $\text{Br}(b \rightarrow s \mu^+ \mu^-)$ branching ratio for the multi-Higgs doublet model normalized by the SM branching ratio in the space of $|Y|^2$ and $\text{Re}(XY^*)$ for $m_H = 100$ GeV. For numerical calculations we use $m_b = 4.7$ GeV and $m_c = 1.5$ GeV. To avoid the large effect of the J/ψ resonance we integrate the differential branching ratio for the kinematical range $4m_\mu^2 < s < (m_{J/\psi} - \delta)^2$ where m_μ is muon mass, $m_{J/\psi}$ is J/ψ mass and $\delta = 100$ MeV. The interference effect between the J/ψ resonance and the short distance contribution in this kinematical region is still sizable ($\lesssim 20\%$) for the SM case [11]. However as we can see in Fig.2 the interference effect is expected to be smaller than the charged Higgs contribution to the short distance part. In order to solve $\text{Im}(XY^*)$ in terms of $|Y|^2$ and $\text{Re}(XY^*)$, we assume $\text{Br}(b \rightarrow s \gamma) = 2.8 \times 10^{-4}$. Since there is a sizable experimental error on this quantity we have just used the SM prediction for the illustration. We expect that the experimental error as well as the theoretical ambiguity will be reduced in future when actual analysis on $b \rightarrow s \ell^+ \ell^-$ is done. Fig.2(b) shows the asymmetry in the same kinematical range. From these two figures we can see that the branching ratio can be a few times larger than that of the SM, which is 3.8×10^{-6} in this kinematical region, and the asymmetry can be as large as 20 % compared to about 5 % in the SM. It is

* The CP violating parameter in the $K^0\text{-}\bar{K}^0$ mixing, ϵ_K , also receives a similar contribution from the charged Higgs loop as the $B^0\text{-}\bar{B}^0$ mixing. In this case, however, the relevant CKM matrix element is different and the constraint on $|Y|$ from ϵ_K only is not strong. If we combine the $B^0\text{-}\bar{B}^0$ mixing and the ϵ_K constraints we can exclude a slightly larger parameter space but the allowed region of $|Y|$ is numerically almost the same as above.

also interesting to see that the branching ratio is sensitive to the value of $|Y|^2$, on the other hand the asymmetry gives an independent constraint on the parameter space. Therefore the values of $|Y|^2$ and $\text{Re}(XY^*)$ are determined if the branching ratio and the asymmetry are measured with reasonable accuracy. Fig.2(c) shows the $b \rightarrow s \nu \bar{\nu}$ branching ratio normalized by the SM prediction ($\text{Br}(b \rightarrow s \nu \bar{\nu})_{SM} = 4.3 \times 10^{-5}$) in the same parameter space. In this case the branching ratio only depends on the parameter $|Y|^2$ and is enhanced by a similar factor as the branching ratio of $b \rightarrow s \ell^+ \ell^-$.

If $\text{Im}(XY^*) \neq 0$, there is possibility to observe the CP violating charge asymmetry, *i.e.* the difference of $b \rightarrow s \ell^+ \ell^-$ and $\bar{b} \rightarrow \bar{s} \ell^+ \ell^-$. Since this quantity is induced with help of the phase in $Y(\hat{s})$ in Eq.(23), the asymmetry appears only above the $c\bar{c}$ threshold of the lepton invariant mass. The charge asymmetry is defined as

$$A_{CP} = \frac{\int_{(m_{\psi'} + \delta)^2}^{m_b^2} ds \left(\frac{d\text{Br}(b \rightarrow s \ell^+ \ell^-)}{ds} - \frac{d\text{Br}(\bar{b} \rightarrow \bar{s} \ell^+ \ell^-)}{ds} \right)}{\int_{(m_{\psi'} + \delta)^2}^{m_b^2} ds \left(\frac{d\text{Br}(b \rightarrow s \ell^+ \ell^-)}{ds} + \frac{d\text{Br}(\bar{b} \rightarrow \bar{s} \ell^+ \ell^-)}{ds} \right)}, \quad (30)$$

where $m_{\psi'}$ is the ψ' mass and we take $\delta = 100$ MeV. In Fig.2(d) this quantity is shown for $m_H = 100$ GeV. Since this asymmetry is at most a few percent, we need large statistics to measure it. Note that there is a twofold ambiguity in sign of $\text{Im}(XY^*)$. Fig.2(a)-(c) do not depend on the sign of $\text{Im}(XY^*)$ while A_{CP} (Fig.2(d)) change its sign according to the sign of $\text{Im}(XY^*)$. Fig.2(d) shows the case of $\text{Im}(XY^*) > 0$.

In Fig.3(a)-(d) we show the same quantities for $m_H = 300$ GeV. As in the case of $m_H = 100$ GeV we can see similar enhancements on the branching ratios and the forward-backward asymmetry in the allowed parameter space.

In summary we have investigated possible constraints on the charged Higgs coupling parameters from the $b \rightarrow s \gamma$, $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ processes. Within the present experimental constraints including $b \rightarrow s \gamma$ process, we have shown that the branching ratio of the $b \rightarrow s \ell^+ \ell^-$ and $b \rightarrow s \nu \bar{\nu}$ processes can be enhanced a few times compared with the SM. The lepton forward-backward asymmetry in the $b \rightarrow s \ell^+ \ell^-$ process can also be enhanced by a factor of several. Since these quantities depend on the model parameters in different ways, we can obtain useful information on the model once these processes are observed experimentally.

This work is supported by the Grant-in-aid for Scientific Research from the Ministry of Education, Science and Culture of Japan.

References


- [1] M.S. Alam *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **74** (1995) 2885.
- [2] T.G. Rizzo, *Phys. Rev.* **D 38** (1988) 820;
W.S. Hou and R.S. Wiley, *Phys. Lett.* **B 202** (1988) 591;
C.Q. Geng and J.N. Ng, *Phys. Rev.* **D 38** (1988) 2857;
V. Barger, J.L. Hewett and R.J.N. Phillips, *Phys. Rev.* **D 41** (1990) 3421.
- [3] T. Goto and Y. Okada, *Prog. Theor. Phys.* **94** (1995) 407, and references therein.
- [4] CLEO Collaboration, R. Balest *et al.*, CLEO-CONF 94-4 (1994);
CDF Collaboration, F. Abe *et al.*, *Phys. Rev. Lett.* **76** (1996) 4675.
- [5] Y. Grossman, Z. Ligeti, and E. Nardi, *Nucl. Phys.* **B 465** (1996) 369;
ALEPH Collaboration, D.Buskulic *et al.*, *Phys. Lett.* **B343** (1995) 444.
- [6] Y. Grossman, *Nucl. Phys.* **B426** (1994) 355.
- [7] Y. Grossman and Y. Nir, *Phys. Lett.* **B313** (1993) 126.
- [8] G.L. Glashow and S. Weinberg, *Phys. Rev.* **D15** (1977) 1958.
- [9] B. Grinstein, M.J. Savage and M.B. Wise, *Nucl. Phys.* **B319** (1989) 271.
- [10] A. Ali, G. Giudice and T. Mannel, *Z. Phys.* **C 67** (1995) 417.
- [11] T. Goto, Y. Okada, Y. Shimizu, and M. Tanaka, hep-ph/9609512, To be published in *Phys. Rev. D*.
- [12] M. Boulware and D. Finnell, *Phys. Rev.* **D44** (1991) 2054.
- [13] LEP Electroweak Working Group and the SLD Heavy Flavor Group, CERN-PPE-96-183(1996).
- [14] ALEPH Collaboration, CERN-PPE-97-018(1997).

FIG. 1: The charged Higgs contribution to δR_b for $m_H = 100$ GeV (solid line) and $m_H = 300$ GeV (dashed line). The dotted line represents the 3σ lower bound from the world average value of R_b .

FIG. 2: (a). The ratio of the $\text{Br}(b \rightarrow s\ell^+\ell^-)$ in the range $4m_\mu^2 \leq s \leq (m_{J/\psi} - \delta)^2$ for the multi-Higgs doublet model (MHDM) and the standard model (SM) in the case of $m_H = 100$ GeV. (b). The forward-backward asymmetry (A_{FB}) in the range $4m_\mu^2 \leq s \leq (m_{J/\psi} - \delta)^2$ for the MHDM and the SM in the case of $m_H = 100$ GeV. (c). The ratio of the $\text{Br}(b \rightarrow s\nu\bar{\nu})$ for the MHDM and the SM in the case of $m_H = 100$ GeV. (d). The charge asymmetry (A_{CP}) in the range $(m_{\psi'} + \delta)^2 \leq s \leq m_b^2$ for $m_H = 100$ GeV. We take $\delta = 100$ MeV in (a), (b) and (d).

FIG. 3: The same as Fig. 2 for $m_H = 300$ GeV.

Figure 1






Figure 2:




Figure 3: